arXiv:1212.5547v1 [physics.soc-ph] 21 Dec 2012

Discontinuous Transition of a Multistage Independent Cascade Model on Networks

Takehisa Hasegawa[1, *] and Koji Nemoto[2, †]

[1] *Graduate School of Information Science, Tohoku University,*
6-3-09, Aramaki-Aza-Aoba, Sendai, Miyagi, 980-8579, Japan
[2] *Department of Physics, Hokkaido University, Kita 10 Nisi 8, Kita-ku, Sapporo, Hokkaido, 060-0810, Japan*

We study a multistage independent cascade (MIC) model in complex networks. This model is parameterized by two probabilities: T_1 is the probability that a node adopting a fad increases the awareness of a neighboring susceptible node until it abandons the fad, and T_2 is the probability that an adopter directly causes a susceptible node to adopt the fad. We formulate a framework of tree approximation for the MIC model on an uncorrelated network with an arbitrary given degree distribution $\{p_k\}$. As an application, we study this model on a random regular network with degree $k = 6$ to show that it has a rich phase diagram including continuous and discontinuous transition lines for the percolation of fads as well as a continuous transition line for the percolation of susceptible nodes. In particular, the percolation transition of fads is discontinuous (continuous) when T_1 is larger (smaller) than a certain value. Furthermore, the phase boundaries drastically change by assigning a finite fraction of initial adopters. We discuss this sensitivity by both tree approximation and numerical simulation.

PACS numbers: 89.75.Hc,87.23.Ge,05.70.Fh,64.60.aq

INTRODUCTION

Word-of-mouth phenomena often catapult books, movies, and music out of obscurity and into popularity. Similarly, we witness similar propagative effects in voting behavior, the spread of rumors, and diffusion of innovations. To explain the mechanism causing such information propagation, Watts proposed a network model (the Watts model) in which the decision made by each node (person) to adopt an innovation is determined by the relative number of its adjacent nodes (friends) who have already adopted it [1]. In this model, the system transits from the global cascade phase, where some small initialized shocks are propagated though a whole system, to the local cascade phase, when the threshold of their decision making increases. This transition is discontinuous: the final fraction of adopters rises discontinuously and its distribution over many trials is bimodal even at the transition threshold. Following the seminal paper by Watts, several authors have investigated the effects of network topology on the transition threshold [2–7].

The Watts model assumes that a node is *permanently active* in the sense that the node will continue to adopt an innovation once it has done so. On the other hand, the susceptible-infected-removed (SIR) model [8] and the independent cascade (IC) model [9] (including the SIR model with transmissibility [10]) have been investigated as models of *transient fads*. In the IC model, an adopter of a fad makes only one attempt, with a certain probability, to persuade each susceptible neighbor to adopt it, and after that attempt, the adopter abandons the fad. The SIR model and the IC model have been extensively studied both analytically and numerically on complex networks [10–12]. On such networks, these models do not show a discontinuous transition, but a continuous one [26].

In recent years, several multistage contagion models have been reported [13–17]. Even when a fad is transient, it may show a discontinuous transition when the number of possible states for each node is increased. Krapivsky et al. proposed a model of transient fads (the fad model) by incorporating *social reinforcement* into the SIR model [16]. Social reinforcement means that each node requires multiple prompts from adjacent adopters before adopting a fad. In this model, a susceptible node takes one of M awareness levels $0, 1, \cdots$, and $M - 1$, and it becomes an adopter when the level reaches M. Each adopter increases the awareness level of the adjacent neighbors by one at unit rate until the adopters begin to abandon the fad at a certain rate. Note that the case of no reinforcement ($M = 1$) coincides with the standard SIR model. Krapivsky et al. studied this model in a mean-field framework to show that the transition is continuous for $M = 1$ and discontinuous for $M \geq 2$. However, nothing is known about how this model behaves when placed on a network.

In this paper, we propose a simple multistage independent cascade (MIC) model, inspired by the fad model. This model is parameterized by two probabilities: T_1 is the probability that a node adopting a fad increases the awareness level of a neighboring susceptible node until the node abandons the fad, and T_2 is the probability in which an adopter directly persuades a susceptible node to adopt the fad. We formulate a framework of tree approximation for the MIC model on an uncorrelated network with an arbitrary given degree distribution $\{p_k\}$. As an application, we study this model on a random regular network with degree $k = 6$ to show that it has a rich phase diagram. The percolation transition of fads is discontinuous (continuous) when T_1 is larger (smaller)



FIG. 1: Transition rules of the MIC model. The values on the lines indicate the transmission probability. The events indicated by the dashed lines ($0 \to 1, 0 \to 2$ and $1 \to 2$) occur when the node is adjacent to an adopter, whereas the event indicated by the solid line ($2 \to 3$) occurs irrespective of the neighbor's state.

than a certain value. Furthermore, the phase boundaries drastically change by setting a finite fraction of initial adopters. We discuss this sensitivity by both tree approximation and numerical simulation.

MODEL

Let us consider a network with N nodes. Each node takes one of the four states: (0) susceptible (without awareness), (1) informed (susceptible with awareness), (2) adopter of the fad, and (3) abandoner of the fad. The dynamics of this model is given as follows (Fig.1):

(i) Select randomly the fraction ρ of nodes as the initial adopters (*seeds*). Set the state of other nodes as being susceptible.

(ii) For each adopter, change each of the neighboring susceptible nodes to be either informed ($0 \to 1$) with probability T_1 or an adopter ($0 \to 2$) with probability T_2 and change each of the neighboring informed nodes to an adopter ($1 \to 2$) with probability $T \equiv T_1 + T_2 (\leq 1)$.

(iii) Change all the old adopters to abandoners ($2 \to 3$).

(iv) Repeat (ii) and (iii) until no adopter exists in the network.

Note that the case with $T_1 = 0$ coincides with the standard IC model (the SIR model with transmissibility T_2), and the case with $T_2 = 0$ is regarded as an IC version of the fad model [27].

In the following sections, we denote the fraction of nodes in states 0, 1, 2, and 3 in the final state by S_0, S_1, S_2, and S_3, respectively. Also, we designate the networks composed of susceptible nodes and abandoners the S_0-network and the S_3-network, respectively. The fractions of the largest components of the S_0-network, and the S_3-network are denoted by $S_0^{\max}$ and $S_3^{\max}$, respectively.

PHASE DIAGRAM OF THE MIC MODEL IN THE $\rho \to 0$ LIMIT

In this section, we consider the dynamics starting from an infinitesimal fraction of seeds ($\rho = 0+$) on uncorrelated networks with a given degree distribution $\{p_k\}$. Newman analyzed the SIR model with a transmissibility T (the MIC model with $T_1 = 0$ and $T_2 = T$) by using generating functions [10, 18]. According to his result, the system experiences two phase transitions in increasing T: the giant component of abandoners appears at the percolation threshold of the S_3-network $T_c = \langle k \rangle / \langle k^2 - k \rangle$ ($\langle \cdot \rangle$ means the average of a quantity weighted by p_k), and the giant component of susceptible nodes disappears at the percolation threshold of the S_0-network $T_{c'}$. Correspondingly, there exist three distinct phases: $S_0^{\max} > 0$ and $S_3^{\max} = 0$ below T_c, $S_0^{\max} > 0$ and $S_3^{\max} > 0$ for $T_c < T < T_{c'}$, and $S_0^{\max} = 0$ and $S_3^{\max} > 0$ above $T_{c'}$. In the following, we generalize his method to the MIC model.

Percolation of the S_3-network

Let u be the probability of a susceptible node not being informed by the prompting of a randomly chosen neighboring (RCN) node, and v be the probability that a RCN node does not adopt the fad in the final state of dynamics. Then u and v are related as

$$u = v + (1 - T)(1 - v) = 1 - T + Tv. \tag{1}$$

To evaluate these values, it is convenient to introduce the generating function of the degree distribution $\{p_k\}$ as

$$F_0(x) = \sum_k p_k x^k, \tag{2}$$

$$F_1(x) = \frac{F_0'(x)}{F_0'(1)} = \frac{1}{\langle k \rangle} \sum_k (k+1) p_{k+1} x^k. \tag{3}$$

By using these functions we can express the probabilities of a node being susceptible and informed, which are equal to S_0 and S_1 respectively, as

$$S_0 = F_0(u), \tag{4}$$

$$S_1 = T_1(1 - v) F_0'(u), \tag{5}$$

$$S_3 = 1 - S_0 - S_1. \tag{6}$$

The last equation follows from the fact that $S_2 = 0$ at the final state. Now let us divide v into two parts, $v = v_0 + v_1$, where v_0 and v_1 are the probabilities of a RCN node being susceptible and informed, respectively. The self-consistent equations for v_0 and v_1 are then given as

$$v_0 = F_1(u), \tag{7}$$

$$v_1 = T_1(1 - v) F_1'(u), \tag{8}$$

or by adding them we have

$$v = F_1(u) + T_1(1-v)F_1'(u), \tag{9}$$

which can be expressed in terms of u as

$$u = 1 - T + TF_1(u) + T_1(1-u)F_1'(u). \tag{10}$$

Let us analyze in some detail the solution of this equation in the following.

For any values of T_1 and T_2, there exists a trivial solution $u = 1$, meaning $S_0 = 1$, $S_1 = 0$, and thus $S_3 = 0$. To find nontrivial solutions we solve (10) with respect to T_2 as

$$T_2 = \varphi_0(u; T_1) \equiv (1-u)\frac{1 + T_1F_1'(u)}{1 - F_1(u)} - T_1. \tag{11}$$

This gives T_2 such that a given value of $u < 1$ is a solution of (10) with a fixed T_1. For small $\bar{u} = 1 - u \ll 1$, (11) is expanded to give

$$\begin{aligned} T_2 = &\frac{1}{c_1} + \frac{c_2(1 - c_1T_1)}{2c_1^2}\bar{u} \\ &+ \frac{(3c_2^2 - 2c_1c_3) + c_1(4c_1c_3 - 3c_2^2)T_1}{12c_1^3}\bar{u}^2 + \cdots , \end{aligned} \tag{12}$$

where $c_n = F_1^{(n)}(1)$. Taking the limit $u \to 1-$ ($\bar{u} \to 0+$), we have

$$T_2 = T_c \equiv c_1^{-1} = \frac{1}{F_1'(1)} = \frac{\langle k \rangle}{\langle k^2 - k \rangle}, \tag{13}$$

above which the trivial solution $u = 1$ becomes unstable and S_3 should be nonzero.

For $T_1 < c_1^{-1} = T_c$ fixed, a nontrivial solution exists only for $T_2 > T_c$ and $S_3 \to 0$ when $T_2 \to T_c+$, so that $T_2 = T_c$ is the continuous transition point of S_3. For $T_1 > T_c$, on the other hand, $u < 1$ is possible even for $T_2 < T_c$. To see this let us rewrite (12) for small $\Delta_1 = T_1 - T_c$ and $\Delta_2 = T_c - T_2$, which has the structure as

$$\Delta_2 = 2a\Delta_1\bar{u} - b\bar{u}^2 + \cdots \tag{14}$$

with some positive constants a and b. For $\Delta_1 > 0$, Δ_2 can be positive for small $\bar{u} > 0$, and thus there are two branches of the nontrivial solution for $T_d^L < T_2 < T_c$ with

$$T_d^L \simeq T_c - \frac{a^2}{b}\Delta_1^2, \tag{15}$$

and the lower branch vanishes at $T_2 = T_c$ while the upper branch survives above T_c giving a positive S_3. For $T_2 < T_d^L$, on the other hand, a nontrivial solution does not exist and the only possibility is $S_3 = 0$. This means that S_3 should jump *discontinuously* from zero to the upper branch at somewhere $T_d^L \le T_2 \le T_d^U = T_c$. In general, T_d^L depends on T_1 and is given by $T_d^L(T_1) = \min_{0<u<1} \varphi_0(u; T_1)$.

Percolation of S_0-network

When T_2 increases with T_1 fixed, the S_0-network gradually shrinks and finally breaks into a lot of finitely connected components at $T_2 = T_{2c'}$. Here, let us examine this percolation problem.

The probabilities that a RCN node of a susceptible node in the S_0-network are susceptible, informed, and an adopter are v_0, v_1, and $(1-T)(1-v)$, respectively. Therefore, the generating function for the degree distribution of the S_0-network is given as

$$\begin{aligned} G_0(x) &= \frac{F_0\left(v_0x + v_1 + (1-T)(1-v)\right)}{F_0(u)} \\ &= \frac{F_0\left(v_0(x-1) + u\right)}{F_0(u)}. \end{aligned} \tag{16}$$

Similarly, the generating function for the excess degree distribution of the S_0-network is derived as

$$G_1(x) = \frac{G_0'(x)}{G_0'(1)} = \frac{F_1\left(v_0(x-1) + u\right)}{F_1(u)}. \tag{17}$$

Because the mean excess degree of the S_0-network is given as $G_1'(1)$, the value of u at the percolation threshold $(T_{1c'}, T_{2c'})$ should satisfy

$$1 = G_1'(1) = v_0\frac{F_1'(u)}{F_1(u)}. \tag{18}$$

In the percolating phase the largest component $S_0^{\max}$ is given by

$$S_0^{\max} = S_0\left(1 - G_0(v_S)\right), \tag{19}$$

where v_S is the probability that an S_0-cluster connected via a randomly chosen link to a susceptible node is finite,

$$v_S = G_1(v_S). \tag{20}$$

Phase diagram for random regular network

As an application, we consider the phase diagram of the MIC model on a random regular network with degree $k = 6$. By applying the present tree approximation, we have the phase diagram in T_1-T_2 space as shown in Fig.2. The transition line of the S_3-percolation is independent of T_1 and located at $T_2 = T_c = 1/(k-1) = 0.2$ (the vertical line). However, the order of transition is continuous if $T_1 < T_c$, and otherwise discontinuous. In Fig.3, we plot S_3 with several values of T_1 as a function of T_2. For large T_1, S_3 is multivalued in $T_d^L < T_2 < T_d^U = T_{2c}$. Since dynamics starts from an infinitesimal fraction of seeds, S_3 remains as the lower branch below T_d^U and jumps to the upper branch at $T_2 = T_d^U$, implying that a discontinuous occurrence of the giant component of the S_3-network at $T_2 = T_d^U$. Such a jump disappears when $T_1 < T_{1c}$.



FIG. 2: Phase diagram of the MIC model with $\rho = 0+$. The thick-solid and thick-dashed lines denote the continuous transition line for percolation of the S_3-network $T_{2c'}$ and of the S_0-network T_{2c}, respectively. The thin-solid and dashed lines denote the upper (T_d^U) and lower (T_d^L) bounds, respectively, between which (10) has two nontrivial solutions.



FIG. 3: Analytical result for S_3 with $T_1 = 0$, 0.2, and 0.4 (from right to left) as a function of T_2. T_d^L and T_d^U indicate the lower and upper bounds, respectively, between which (10) has two nontrivial solutions.

The region of coexistence of the giant components of the S_0- and S_3-networks exists in $T_{2c} < T_2 < T_{2c'}(T_1)$ for small T_1. This region shrinks with increasing T_1, and $T_{2c'}(T_1)$ reaches T_{2c} at a certain value of T_1 above which the percolation of the S_3-network and the disappearance of the giant component of the S_0-network occur simultaneously at $T_2 = T_{2c}$.

The tricritical point (T_{1c}, T_{2c}) of the random regular network with degree k is given by $(1/(k-1), 1/(k-1))$. We note that $(T_{1c}, T_{2c}) = (0.5, 0.5)$ when $k = 3$, so there is no room for $T_1 > T_{c1}$ at $T_2 = T_{2c}$ since $T = T_1 + T_2 \leq 1$. This shows that the MIC model on the random regular network with degree $k = 3$ does not show the discontinuous transition of the S_3-network, but only the continuous one.



FIG. 4: Phase diagram of the MIC model with $\rho = 0.01$. The thin-solid and dashed lines denote the upper (T_d^U) and lower (T_d^L) bounds, respectively, between which (10) has two nontrivial solutions. The thick-dashed line denotes the transition line of percolation of the S_0-network $T_{2c'}$. The marks denote the transition point of percolation of the S_3-network obtained by Monte-Carlo simulations.

SENSITIVITY TO THE FRACTION OF INITIAL ADOPTERS

Tree approximation

On the basis of the formalism in the previous section, it is easy to find the self-consistent equations for the case where the fraction of initial adopters is $\rho > 0$. Let $\bar{\rho} = 1 - \rho$ be the initial fraction of susceptible nodes. Then for the present model $v = v_0 + v_1$ where

$$v_0 = \bar{\rho} F_1(u), \quad v_1 = \bar{\rho} T_1 (1 - v) F_1'(u), \tag{21}$$

the relation $u = 1 - T + Tv$ is unchanged. In terms of u, we thus obtain

$$u = 1 - T\left(1 - \bar{\rho} F_1(u)\right) + \bar{\rho} T_1 (1 - u) F_1'(u). \tag{22}$$

By using the solution, we obtain

$$S_0 = \bar{\rho} F_0(u), \quad S_1 = \bar{\rho} \frac{T_1(1-u)}{T} F_0'(u), \tag{23}$$

and $S_3 = 1 - S_0 - S_1$.

For $\rho > 0$, percolation transition of the S_3-network, from zero to nonzero $S_3^{\max}$, can take place as long as ρ is smaller than the percolation density (see the next subsection). In contrast, there is no trivial solution of (22) for all regions of parameters and no continuous transition from zero to nonzero S_3 because all seeds will become abandoners, so $S_3 \geq \rho$. Nevertheless, discontinuous transition of S_3 is still possible. Equation (11) is modified as

$$T_2 = \varphi_\rho(u; T_1) = (1 - u) \frac{1 + \bar{\rho} T_1 F_1'(u)}{1 - \bar{\rho} F_1(u)} - T_1, \tag{24}$$



FIG. 5: Comparison of S_3 with the numerical result for $T_1 = 0.4$ and $\rho = 0.01$. The analytic result (solid curve) and the result obtained by simulation with $N = 1000(\times)$, $2000(+)$, $4000(\square)$, $8000(\blacksquare)$, and $16000(\bigcirc)$ are shown. Here $S_3/(1 - S_3)$ is plotted instead of S_3 itself to see the behaviors near $S_3 = 0$ and 1. The analytic evaluation gives $T_d^L \simeq 0.1306$ and $T_d^U \simeq 0.1587$.



FIG. 6: Monte Carlo results for $S_3^{\max}$ with $T_1 = 0.4$ and $\rho = 0.01$ as a function of T_2. Data for $N = 1000(\times)$, $2000(+)$, $4000(\square)$, $8000(\blacksquare)$, and $16000(\bigcirc)$ are shown. The discontinuous transition point of S_3 and the transition point of S_3-percolation are estimated from these data as $T_2 \sim 0.158$ and $T_2 \sim 0.106$, respectively.

and the lower bound T_d^L and the upper bound T_d^U of the discontinuous transition point can be given by the local minimum and local maximum of $\varphi_\rho(u; T_1)$ in $0 < u < 1$, respectively. We note that there is a value of ρ, depending on T_1, above which $\varphi_\rho(u; T_1)$ is monotonically decreasing and these bounds do not exist (see Fig.8).

For the S_0-network, the formulation for deriving the percolation threshold $T_{c'}$ and the ratio of the largest component $S_0^{\max}$ is unchanged, so that (18) and (19) still hold with (21) and (23).



FIG. 7: Monte Carlo results for $S_0^{\max}$ with $T_1 = 0.25$ as a function of T_2. Data for $N = 1000(\times)$, $2000(+)$, $4000(\square)$, $8000(\blacksquare)$, and $16000(\bigcirc)$ are shown. The solid line is drawn by (19), and gives $T_{2c'} \simeq 0.22576$.

Results for random regular network

Now let us discuss the sensitivity of the fraction of the initial adopters by considering the random regular network with degree 6. Here, we performed Monte Carlo simulations of the MIC model to verify our analytical results. The number of nodes used is $N = 1000$, 2000, 4000, 8000, and 16000. The number of graph realizations is 100, and the number of trials on each graph is 100.

In Fig.4, we show the phase diagram for the case with $\rho = 0.01$. The upper and lower bounds of the discontinuous transition of the S_3-network and the percolation threshold of the S_0-network are drawn by the tree approximation. We find that the upper bound shifts to the left from the vertical line (the discontinuous transition line of the case with $\rho = 0+$). We also plot numerical results for the percolation transition points of the S_3-network (our tree approximation does not give $S_3^{\max}$ when $\rho > 0$). Each transition point is estimated from the cross point of the fractal exponent [19, 20], which is defined as

$$\begin{aligned}\psi(N) &\equiv \lim_{N\to\infty} \log_N S_3^{\max}(N) \\ &\approx \frac{\log S_3^{\max}(2N) - \log S_3^{\max}(N/2)}{\log 2N - \log N/2}. \qquad (25)\end{aligned}$$

Note that the crossing point of the fractal exponent indicates that the transition is continuous. We find that the continuous transition points of the S_3 network are outside the interval between the lower and upper bounds. This means that the percolation of the S_3-network and the discontinuous jump of S_3 occur at different points: when T_2 increases with T_1 fixed, the continuous transition of the S_3-network occurs first, and then a discontinuous change in S_3 occurs at a larger value of T_2 between the lower and upper bounds.



FIG. 8: T_2 dependence of S_3 with $T_1 = 0.4$ and $\rho = 0$, 0.01, and 0.1 (from left to right).

In Fig.5, we compare the analytical result and numerical evaluations for S_3 with $T_1 = 0.4$. We find that the analytical result coincides well with the numerical one except near the discontinuous transition point $T_2 = T_d^U(T_1) \simeq 0.1587$. Near the transition point, numerically evaluated S_3 approaches the analytical curve when N increases. Although it is not analytically confirmed that the upper bound T_d^U coincides with the actual discontinuous transition point when the fraction of the initial adopters is finite, the jump of S_3 from the lower branch to the upper branch occurs very near or just at T_d^U.

We plot the Monte Carlo result for $S_3^{\max}$ with $T_1 = 0.4$ and $S_0^{\max}$ with $T_1 = 0.25$ for $\rho = 0.01$ in Figs.6 and 7, respectively. From Fig.6 we find that the S_3-network first percolates but grows discontinuously at a larger value of T_2. In Fig.7, we have a good correspondence between the numerical results for $S_0^{\max}$ and Eq.(19).

Finally, we show analytical curves of S_3 with several values of ρ and a fixed $T_1 = 0.4$ in Fig.8. When ρ is small, we observe the precursor of the discontinuous transition of S_3 in the same way as in Fig.3. However, such a behavior disappears with increasing ρ. The system does not show the discontinuous transition if the fraction of the initial adopters ρ is sufficiently large.

SUMMARY

We have investigated a simple multistage independent cascade (MIC) model on networks. Under the local tree approximation, we have derived the formulation for evaluating S_0, S_1, and S_3 and thus the phase diagram. In particular, percolation of the adopters occurs discontinuously for large values of T_1. Moreover, when dynamics starts from a finite fraction of initial adopters, the discontinuous jump in the number of adopters occurs after the adopters percolate. As an application, we have studied the phase diagram of the MIC model on a random regular graph and confirmed that the results support well our analytical predictions.

Our tree approximation for the present MIC model is valid only for uncorrelated networks, and it is beyond our scope to investigate how the cooperative behavior of this model depends on the properties of the network on which the dynamics is performed (e.g., clustering, degree correlation, and community structure). For example, clustering decreases the threshold of the SIR model [21], but an innovation in the Watts model is easier to propagate in rather clustered networks (as compared to random networks) [2, 6]. How does the MIC model behave in clustered networks? In order to answer such questions, further research is needed.

Acknowledgements

TH acknowledges the support provided by the Japan Society for the Promotion of Science through Grant-in-Aid for Young Scientists (B) (Grant No. 24740054).

---

[*] Electronic address: hasegawa@m.tohoku.ac.jp
[†] Electronic address: nemoto@statphys.sci.hokudai.ac.jp

[1] D. J. Watts, Proc. Natl. Acad. Sci. USA **99**, 5766 (2002).
[2] D. Centola, V. M. Eguíluz, and M. W. Macy, Physica A **374**, 449 (2007).
[3] J. P. Gleeson and D. J. Cahalane, Phys. Rev. E **75**, 056103 (2007).
[4] J. P. Gleeson, Phys. Rev. E **77**, 046117 (2008).
[5] J. L. Payne, P. S. Dodds, and M. J. Eppstein, Phys. Rev. E **80**, 026125 (2009).
[6] Y. Ikeda, T. Hasegawa, and K. Nemoto, in *J. Phys. Conf. Ser.* (2010), vol. 221, p. 012005.
[7] A. Hackett, S. Melnik, and J. P. Gleeson, Phys. Rev. E **83**, 056107 (2011).
[8] R. M. Anderson and R. M. May, *Infectious Diseases of Humans: Dynamics and Control* (Oxford University press, 1992).
[9] D. Kempe, J. Kleinberg, and É. Tardos, in *Proceedings of the ninth ACM SIGKDD international conference on Knowledge discovery and data mining* (2003), pp. 137–146.
[10] M. E. J. Newman, Phys. Rev. E **66**, 016128 (2002).
[11] R. Pastor-Satorras and A. Vespignani, Phys. Rev. Lett. **86**, 3200 (2001).
[12] R. Pastor-Satorras and A. Vespignani, Phys. Rev. E **63**, 066117 (2001).
[13] T. Antal and P. L. Krapivsky, J. Stat. Mech. Theor. Exp. **2012**, P07018 (2012).
[14] C. de Kerchove, G. Krings, R. Lambiotte, P. V. Dooren, and V. D. Blondel, Phys. Rev. E **79**, 016114 (2009).
[15] S. Melnik, J. Ward, J. Gleeson, and M. Porter, arXiv: 1111.1596 (2011).
[16] P. L. Krapivsky, S. Redner, and D. Volovik, J. Stat. Mech. Theor. Exp. **2011**, P12003 (2011).

[17] M. Starnini, A. Baronchelli, and R. Pastor-Satorras, J. Stat. Mech. Theor. Exp. **2012**, P10027 (2012).

[18] M. E. J. Newman, Phys. Rev. Lett. **95**, 108701 (2005).

[19] T. Nogawa and T. Hasegawa, J. Phys. A: Math. Theor. **42**, 145001 (2009).

[20] T. Hasegawa, T. Nogawa, and K. Nemoto, arXiv: 1009.6009 (2010).

[21] M. E. J. Newman, Phys. Rev. E **68**, 026121 (2003).

[22] T. Tomé and R. M. Ziff, Phys. Rev. E **82**, 051921 (2010).

[23] E. Kenah and J. M. Robins, Phys. Rev. E **76**, 036113 (2007).

[24] J. C. Miller, Phys. Rev. E **76**, 010101 (2007).

[25] E. Kenah and J. C. Miller, Interdiscip. Perspect. Infect. Dis. **2011** (2011).

[26] The SIR model is mapped onto the IC model with a transmission probability (the SIR model with transmissibility). Although these two behave differently from each other on local scales, there is a correspondence between these two for critical points and critical exponents [22–25].

[27] The adopters of the original fad model can infect a susceptible neighbor twice or more, while those in the present model can infect each susceptible neighbor at most once.